

effective for dissolving fatty acids having a chain length of an odd number of carbon atoms and derivatives thereof (refer to, for example, patent document 3).

On the other hand, it is known as well that
5 polyhydric alcohols are effective as a solubilizing agent _____
for 6-benzylaminopurine (refer to, for example, patent document 4).

However, it is difficult to stabilize dissolution of a preparation in which 6-benzylaminopurine and
10 derivatives thereof are coexistent with fatty acids having a chain length of an odd number of carbon atoms and derivatives thereof, and in particular, the problem that active ingredients in the preparation are deposited at low temperature is involved therein. Accordingly, the
15 stability at low temperature is not yet satisfactory in terms of actual use, and the existing situation is that hair growth promotor compositions having further excellent dissolution stability are still desired.

Patent document 1: Japanese Patent Application Laid-Open
20 No. 320028/1993 (claims, examples and other)

Patent document 2: Japanese Patent Application Laid-Open
No. 233037/1995 (claims, examples and other)

Patent document 3: Japanese Patent publication No.
37886/1990 (claims, examples and other)

25 Patent document 4: Japanese Patent Application Laid-Open

No. 72321/1998 (claims, examples and other)

Disclosure of the Invention


In light of the problems of the prior art described
5 ~~above and the existing situation, the present invention~~
intends to solve them, and an object thereof is to
provide a hair growth promotor composition which provides
an excellent hair growth effect by applying externally
onto a scalp and which is excellent particularly in
10 stability at low temperature and has a good use feeling.

Intensive researches repeated by the present
inventors in order to solve the conventional problems
described above have resulted in finding that a hair
growth promotor composition which meets the object
15 described above and which is excellent in stability at
low temperature is obtained when containing at least one
compound selected from fatty acids having a chain length
of an odd number of carbon atoms, the derivatives of the
fatty acids, aliphatic alcohols having a chain length of
20 an odd number of carbon atoms and the derivatives of the
aliphatic alcohols, 6-benzylaminopurine and/or a
derivative thereof and two kinds of fatty acid esters,
and thus the present invention has come to be completed.

That is, the hair growth promotor composition of
25 the present invention is an ethanol or aqueous ethanol

preparation comprising, as active ingredients for a hair growth promotor, (A) at least one compound selected from fatty acids having a chain length of an odd number of carbon atoms, the derivatives of said fatty acids,

5 aliphatic alcohols having a chain length of an odd number of carbon atoms and the derivatives of said aliphatic alcohols and (B) at least one selected from 6-benzylaminopurine and/or the derivatives thereof represented by the following Formula (I), wherein it
10 further comprises (C) at least one of polyglycerin fatty acid esters and (D) at least one of sorbitan fatty acid esters:



in Formula (I), R₁ is an alkyl group having 1 to 22 carbon atoms, a cyclic hydrocarbon group, an alkenyl
15 group having 1 to 22 carbon atoms, an aralkyl group which is non-substituted or has a substituent, a styryl group which is non-substituted or has a substituent, an alkylamino group, an amino group having a cyclic hydrocarbon group, an alkenylamino group, a benzylamino
20 group which is non-substituted or has a substituent, a

phenylethylamino group which is non-substituted or has a substituent, a phenylamino group which is non-substituted or has a substituent, a phenylaminocarbonylamino group which is non-substituted or has a substituent, a

5 pyridylamino group, a pyridylmethylamino group, a pyrrolylmethylamino group, an oxazolylmethylamino group, an imidazolylmethylamino group, a pyridazolylmethylamino group, a naphthylamino group or a naphthylmethylamino group, and R₂ is a hydrogen atom, a pentose or hexose.

10

Effects of the Invention

According to the present invention, provided is a hair growth promotor composition which has an excellent hair growing effect and provides an excellent

15 stabilization effect at low temperature and which provides a good use feeling having no stickiness.

Best Mode for Carrying Out the Invention


The embodiment of the present invention shall be
20 explained below in details.

The hair growth promotor composition of the present invention is an ethanol or aqueous ethanol preparation comprising, as active ingredients for a hair growth promotor, (A) at least one compound selected from fatty
25 acids having a chain length of an odd number of carbon

atoms, the derivatives of said fatty acids, aliphatic alcohols having a chain length of an odd number of carbon atoms and the derivatives of said aliphatic alcohols and (B) at least one selected from 6-benzylaminopurine and/or

5 the derivatives thereof represented by the following

Formula (I), wherein it further comprises (C) at least one of polyglycerin fatty acid esters and (D) at least one of sorbitan fatty acid esters:



in Formula (I), R_1 is an alkyl group having 1 to 22 carbon atoms, a cyclic hydrocarbon group, an alkenyl group having 1 to 22 carbon atoms, an aralkyl group which is non-substituted or has a substituent, a styryl group which is non-substituted or has a substituent, an alkylamino group, an amino group having a cyclic hydrocarbon group, an alkenylamino group, a benzylamino group which is non-substituted or has a substituent, a phenylethylamino group which is non-substituted or has a substituent, a phenylamino group which is non-substituted or has a substituent, a phenylaminocarbonylamino group

which is non-substituted or has a substituent, a pyridylamino group, a pyridylmethylamino group, a pyrrolylmethylamino group, an oxazolylmethylamino group, an imidazolylmethylamino group, a pyridazolylmethylamino group, a naphthylamino group or a naphthylmethylamino group, and R_2 is a hydrogen atom, a pentose or hexose.


In the present invention, the fatty acid having a chain length of an odd number of carbon atoms which is used as the component (A) may be saturated fatty acid or unsaturated fatty acid as long as it has an odd number of carbon atoms constituting the carbon chain and has a hair growth action.

The unsaturated fatty acid may contain plural double bonds. The carbon chain has preferably 9 to 29 carbon atoms, more preferably 11 to 25 carbon atoms, and the carbon chain may be linear or branched. To be more specific, capable of being given as such a fatty acid are, for example, nonanoic acid, undecanoic acid, tridecanoic acid, pentadecanoic acid, nonadecanoic acid, heneicosanoic acid, tricosanoic acid, pentacosanoic acid and heptacosanoic acid.


Any derivatives of the fatty acids having a chain length of an odd number of carbon atoms described above can be used as the usable derivatives of the fatty acids having a chain length of an odd number of carbon atoms as

long as they contain fatty acid residues of fatty acids having an odd number of carbon atoms and can be used for human bodies. The particularly preferred derivatives include the following compounds of (a) to (m).

5 (a) Monoglyceride represented by the following Formula (II) or (III):




..... (II)




..... (III)

In Formula (II) or (III), R represents a linear or branched aliphatic hydrocarbon group having a chain
10 length of an even number of carbon atoms.

(b) Diglyceride represented by the following Formula (IV) or (V):



..... (IV)



..... (V)

In Formula (IV) or (V), R_1 and R_2 are a linear or
15 branched aliphatic hydrocarbon group, and at least one of them represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms. If either one of R_1 and R_2 is a linear

or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms, effects of the present invention are obtained, and the other may be a linear or branched aliphatic hydrocarbon group having a

5 chain length of an odd number of carbon atoms. Further preferably, both of R_1 and R_2 are linear or branched aliphatic hydrocarbon groups having a chain length of an even number of carbon atoms.

(c) Triglyceride represented by the following Formula

10 (VI):



In Formula (VI), R_1 , R_2 and R_3 are a linear or branched aliphatic hydrocarbon group, and at least one of them represents a linear or branched aliphatic

15 hydrocarbon group having a chain length of an even number of carbon atoms. If at least one of R_1 , R_2 and R_3 is a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms, effects of the present invention are obtained, and the others may

20 be linear or branched aliphatic hydrocarbon groups having a chain length of an odd number of carbon atoms.

Preferably, two or more of R_1 , R_2 and R_3 are linear or branched aliphatic hydrocarbon groups having a chain length of an even number of carbon atoms. Further preferably, all of R_1 , R_2 and R_3 are linear or branched

5 ~~aliphatic hydrocarbon groups having a chain length of an~~
even number of carbon atoms.

(d) Fatty acid salt represented by the following Formula (VII):




10 In Formula (VII), R represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms. M represents a metal atom or an ammonium ion. Further, n represents an integer corresponding to a valency of M.

15 (e) Ester represented by the following Formula (VIII):




In Formula (VIII), R represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms. R' represents
20 a monohydric or dihydric aliphatic alcohol residue having an even number of carbon atoms, a polyoxyethylene residue, a sorbitan residue or a sucrose residue.

(f) Primary amide represented by the following Formula (IX):




In Formula (IX), R represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms. R' and R'' represent hydrogen, an alkyl group or a hydroxyalkyl group.

5 (g) Secondary amide represented by the following Formula (X):



In Formula (X), R₁ and R₂ are an aliphatic hydrocarbon group, and either one of them represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms. R' represents hydrogen, an alkyl group or a hydroxyalkyl group.

(h) Tertiary amide represented by the following Formula (XI):




15

In Formula (XI), R₁, R₂ and R₃ are a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms, and at least one of them represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number

20


of a carbon atoms.

(i) Dibasic acid represented by the following Formula (XII) and salt thereof:



5 In Formula (XII), R represents a linear or branched aliphatic hydrocarbon group having a chain length of an odd number of carbon atoms.

(j) Sterol ester represented by the following Formula (XIII):




10

In Formula (XIII), R represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms.

(k) Phospholipid represented by the following Formula


15 (XIV):



In Formula (XIV), R_1 and R_2 are an aliphatic


hydrocarbon group, and either one of them represents a linear or branched aliphatic hydrocarbon group having a chain length of an odd number of carbon atoms. X represents a choline residue, an ethanolamine residue, a serine residue or an inositol residue.

(l) Phosphatidic acid represented by the following Formula (XV):



In Formula (XV), R_1 and R_2 are an aliphatic hydrocarbon group, and either one of them represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms.

(m) Sphingolipid represented by the following Formula (XVI):



In Formula (XVI), R represents a linear or branched aliphatic hydrocarbon group having a chain length of an even number of carbon atoms. X represents a saccharide residue, a phosphoric acid residue or an amine base

residue.

In the present invention, the aliphatic alcohol of the component (A) which is preferably used as the hair growth component may have either saturated or unsaturated carbon chain as long as the number of carbon atoms constituting the carbon chain is an odd number. When it is unsaturated, it may contain plural double bonds. Further, the alcohol may be lower alcohol or higher alcohol and may be any of primary, secondary and tertiary alcohols.

Capable of being given as the preferred aliphatic alcohol having an odd number of carbon atoms are, for example, n-propyl alcohol, n-amyl alcohol, n-heptyl alcohol, n-nonyl alcohol, n-undecyl alcohol, n-tridecyl alcohol, n-pentadecyl alcohol, n-heptadecyl alcohol, n-nonadecyl alcohol, n-heneicosyl alcohol, n-tricosyl alcohol and n-pentacosyl alcohol.

Further, the derivatives of the above aliphatic alcohols having an odd number of carbon atoms can be used as well in the present invention. The representative derivatives thereof are esters and ethers of alcohols having an odd number of carbon atoms.

The preferred esters include esters of the aliphatic alcohols having an odd number of carbon atoms

described above with aliphatic carboxylic acids
(particularly preferably carboxylic acids having 2 to 24
carbon atoms), organic acids such as succinic acid,
citric acid, fumaric acid, lactic acid, pyruvic acid,
-----5-----malic acid and oxalacetic acid and inorganic acids such
as phosphoric acid.

The preferred ethers include ethers of the
aliphatic alcohols having an odd number of carbon atoms
described above with aliphatic alcohols (particularly
10 preferably alcohols having 2 to 24 carbon atoms),
polyhydric alcohols such as glycerin, polyglycerin,
ethylene glycol, propylene glycol and butanediol and
saccharides such as fructose, ribose, galactose,
arabinose, mannose, xylose, sorbitol and mannitol.

15 Further, the ethers described above may contain two
or more aliphatic alcohol residues having an odd number
of carbon atoms in a molecule as is the case with, for
example, di- or tri- alkoxides of alcohols having an odd
number of carbon atoms of glycerin. The derivatives of
20 the aliphatic alcohols having an odd number of carbon
atoms used for the composition of the present invention
may only contain the alcohol residues having an odd
number of carbon atoms described above as long as they do
not exert adverse effects on human bodies. Accordingly,
25 acid residues in the esters described above and alcohol

residues and saccharide residues in the ethers may be substituted with various substituents.

The component (A) used in the present invention is preferably glycerin monoesters of fatty acids having an odd number of carbon atoms and includes, to be specific, glyceride of pentadecanoic acid, glyceride of tridecanoic acid and glyceride of heptadecanoic acid, and they provide an excellent hair growth effect.

The compounds selected from the above fatty acids having a chain length of an odd number of carbon atoms or the derivatives thereof and the aliphatic alcohols having a chain length of an odd number of carbon atoms or the derivatives thereof can be used, if necessary, alone or in combination of two or more kinds thereof, and the content thereof is preferably, 0.001 to 20 mass % (hereinafter referred to only as %), particularly preferably, 0.1 to 10 % based on the total amount of the hair growth promotor composition.

If the above content of the component (A) is less than 0.001 %, the satisfactory hair growth effect is not exhibited in a certain case. On the other hand, if it exceeds 20 %, further more effects are not usually exhibited.

In Formula (I) representing the component (B) used in the present invention, the group represented by R_1

includes the aforementioned respective group, and it is preferably a benzyl group which is non-substituted or has a substituent from the viewpoint of further exhibiting the effects of the present invention. To be specific, it


- 5 ~~includes a benzyl group, a 2-methylbenzyl group,~~
- a 3-methylbenzyl group, a 4-methylbenzyl group,
 a 4-ethylbenzyl group, a 3-chlorobenzyl group,
 a 4-chlorobenzyl group, a 2,4-dichlorobenzyl group,
 a 2-hydroxybenzyl group, a 3-fluorobenzyl group,
 10 a 4-nitrobenzyl group, a 4-bromobenzyl group,
 a 4-fluorobenzyl group, a 3-nitrobenzyl group,
 a 4-propylbenzyl group, a 3,5-difluorobenzyl group,
 a 2-cyanobenzyl group, a 2-acetaminobenzyl group,
 a 4-acetaminobenzyl group,
 15 a 4-methoxycarbonylbenzyl group,
 a 4-dimethylaminobenzyl group, a 4-methoxybenzyl group,
 a 3-trimethylsilyloxybenzyl group,
 a 3-trifluoromethylbenzyl group,
 a 4-butyldimethylsilyloxybenzyl group,
 20 a 2-methoxybenzyl group,
 a 4-trimethylsilyloxybenzyl group and
 a 4-methylthiobenzyl group.

The group represented by R_2 includes preferably, from the viewpoint of further exhibiting the effects of
 25 the present invention, a hydrogen atom, pentose (it

includes, for example, a 1-ribofuranosyl group, a 1-lyxofuranosyl group, a 1-xylofuranosyl group and a 1-arabofuranosyl group) and hexose (it includes, for example, a 1-glucosyl group, a 1-galactosyl group, a 1-gulose group, a 1-mannosyl group and a 1-allose group.


The component (B) used in the present invention is the substance represented by Formula (I) described above, and both of the substances obtained by refining a natural product according to an ordinary method and the substance
10 obtained by synthesis can be used.

The specific compound represented by Formula (I) described above which can preferably be used includes 6-benzylaminopurine and the derivatives thereof which are shown below.




15






6-benzylaminopurine



6-(4-methylbenzylamino)purine



6-(4-chlorobenzylamino)purine

6-(4-methoxybenzyl-
amino)purine5 6-(4-trifluoromethylbenzyl-
amino)purine6-(4-methylsulfonyl-
benzylamino)purine6-benzylamino-9-
ribofuranosylpurine6-benzylamino-9-
glucosylpurine

In the present invention, 6-benzylaminopurine represented by Formula (I) and the derivatives thereof can suitably be used alone or in combination of two or
5 more kinds thereof, and they can be added at an optional----- concentration.

The component (B) is particularly preferably 6-benzylaminopurine and 6-(4-methylbenzylamino)purine from the viewpoint of exhibiting an excellent hair growth
10 effect.

A content of the component (B) used in the present invention is preferably, 0.0001 to 10 %, more preferably, 0.01 to 5 % based on the total amount of the hair growth promotor composition.

15 If the content of the component (B) is less than 0.0001 %, the hair growth effect is not sufficiently exhibited in a certain case. On the other hand, if it exceeds 10 %, further more effects are not usually exhibited.

20 In the present invention, the polyglycerin fatty acid ester used as the component (C) shall not be restricted by a polymerization degree of glycerin, a polymerization position of glycerin, a carbon chain length, the presence of unsaturation in the carbon chain,
25 linearity or branching of the carbon chain, the presence

of a functional group in the carbon chain and mono-, di-, tri-, or poly-ester. To be specific, it includes triglycerin monolaurate, triglycerin dilaurate, triglycerin monomyristate, triglycerin isostearate, ~~5 tetraglycerin monooleate, tetraglycerin monocaprate,~~ pentaglycerin monomyristate, pentaglycerin dipalmitate, pentaglycerin monotridecanoate, hexaglycerin diisostearate, hexaglycerin monocondensed recinoleate, hexaglycerin trilaurate, hexaglycerin tetralaurate, 10 heptaglycerin monocaprylate, heptaglycerin trilaurate, heptaglycerin dimyristate, heptaglycerin tristearate, heptaglycerin tetraoleate, decaglycerin monomyristate, decaglycerin monocondensed recinoleate, decaglycerin diisostearate, decaglycerin tetrapentadecanoate, 15 decaglycerin hexaoleate, decaglycerin heptastearate and decaglycerin monorecinoleate, and they can be used alone or in a mixture of two or more kinds thereof.

Among the above polyglycerin fatty acid esters, the esters having a polymerization degree of 3 or more of 20 glycerin are preferred, and the esters having a polymerization degree of 3 to 10 are particularly preferred. If the polymerization degree is 2 or less, the targeted effects are not obtained in a certain case. On the other hand, if the polymerization degree is 11 or 25 more, the raw material is not of a general purpose, and

therefore it is not economical in a certain case. To be specific, pentaglycerin monomyristate, decaglycerin monomyristate and pentaglycerin monooleate are suitable.

A content of the polyglycerin fatty acid esters of
~~5 the component (C) can suitably be controlled according to~~
the contents of the component (A) and the component (B) each described above, and it is preferably 0.001 to 10 %, particularly preferably, 0.01 to 5 % based on the total amount of the hair growth promotor composition.

10 If the content of the component (C) is less than 0.001 %, the targeted stabilization effect at low temperature is not sufficiently exhibited in a certain case. On the other hand, if it exceeds 10 %, further more effects are not usually exhibited.

15 In the present invention, the sorbitan fatty acid ester used as the component (D) shall not be restricted by a carbon chain length, the presence of unsaturation in the carbon chain, linearity or branching of the carbon chain and mono-, di-, tri-, or poly-ester, and it
20 includes, to be specific, sorbitan monolaurate, sorbitan monostearate, sorbitan tristearate, sorbitan monoisostearate, sorbitan monooleate and sorbitan sesquioleate. They can be used alone or in a mixture of two or more kinds thereof.

25 Among the above components (D), the fatty acid

monoesters of sorbitan are preferred and include, to be specific, sorbitan coconut oil fatty acid ester and sorbitan monolaurate.

A content of the sorbitan fatty acid esters of the component (D) is preferably 0.001 to 10 %, particularly preferably, 0.01 to 5 % based on the total amount of the hair growth promotor composition.

If the content of the component (D) is less than 0.001 %, the stabilization effect at low temperature is not sufficiently exhibited in a certain case. On the other hand, if it exceeds 10 %, further more effects are not usually exhibited, and an adverse effect such as stickiness is exerted on the use feeling.

In the present invention, the hair growth promotor composition which contains, as the essential components, a compound of the component (A) described above and at least one selected from 6-benzylaminopurine and/or the derivatives thereof represented by Formula (I) described above, which is the component (B) and which are further contains the polyglycerin fatty acid ester of the component (C) described above and the sorbitan fatty acid ester of the component (D) described above, whereby obtained is the hair growth promotor composition which exhibits a particularly excellent hair growth effect and in which active ingredients are stable over a long period

of time and practicality is high, and the essential components can be added at optional concentrations.

Further, the content proportions in which a synergistic effect of the constitutional substances

5 ~~described above contained in the hair growth promotor~~ composition is exerted to a maximum extent have been investigated to result in finding that a mass ratio of component (C)/ component (D) is preferably 10/1 to 1/60, more preferably 5/1 to 1/10.

10 The targeted stabilization effect at low temperature can be exhibited to a maximum extent by controlling a mass ratio of component (C)/ component (D) to 10/1 to 1/60.

Since the hair growth promotor composition of the
15 present invention is an ethanol or aqueous ethanol preparation, it can contain, in addition to the component (A) to the component (D) each described above, ethanol (fermented ethanol, synthesized ethanol, denatured ethanol and the like) and water (refined water, distilled
20 water, ion-exchanged water, purified water, ultrapure water and the like) as the balance. In addition to the respective components described above, it may contain optional components according to the use purposes thereof. Capable of being given as such components are, for
25 example, nonionic surfactants, glucide base surfactants,

other surfactants, celluloses, oils & fats, ester oils, high molecular resins, coloring agents, fragrances, UV absorbers and medicinal components such as vitamins, hormones, vasodilator agents, amino acids,

5 antiinflammatory agents, cutaneous function accelerators and keratolytic drugs.

The celluloses include hydroxymethyl cellulose, hydroxypropyl cellulose and hydroxypropylmethyl cellulose; the surfactants include polyoxyethylene-
 10 hardened castor oil, polyoxyethylene-hardened castor oil mono- or isostearate and glycerin fatty acid esters; the oils & fats include polyhydric alcohol fatty acid esters (glycerin tri-2-ethylhexanoate, trimethylpropane triisostearate, and the like), safflower oil, evening
 15 primrose oil and jojoba oil; the ester oils include unsaturated fatty acid alkyl esters (ethyl oleate, isopropyl linoleate and the like), methyl myristate and isopropyl myristate; the amino acids include methionine, serine, glycine and cystine; the keratolytic drugs
 20 include salicylic acid and resorcin; the high molecular resins include amphoteric, cationic, anionic and nonionic polymers; and the UV absorbers include octyl methoxycinnamate (Neoheliopan AV), oxybenzone and urocanic acid.

25 The hair growth promotor composition of the present

invention can be used as an external preparation in the form of a homogeneous solution, a lotion and a gel. The hair growth promotor composition of the present invention can assume the form of an aerosol composition, and in
5 such case, it can contain, in addition to the components described above, lower alcohols such as n-propyl alcohol and isopropyl alcohol, combustible gases such as butane, propane, isobutane, liquefied petroleum gas and dimethyl ether and compressed gases such as nitrogen gas, oxygen
10 gas, carbon dioxide gas and nitrous oxide gas.

In the hair growth promotor composition of the present invention thus constituted, an excellent hair growth effect is obtained by combined use of at least one compound selected from the fatty acids having a chain
15 length of an odd number of carbon atoms, the derivatives of said fatty acids, the aliphatic alcohols having a chain length of an odd number of carbon atoms and the derivatives of said aliphatic alcohols with 6-benzylaminopurine and/or the derivatives thereof, and
20 reduction in stability at low temperature and use feeling caused by the combined use described above are improved without damaging the hair growth effect described above by further adding at least one of polyglycerin fatty acid esters and at least one of sorbitan fatty acid esters.

EXAMPLES

Next, the present invention shall be explained in further detail with reference to examples and comparative examples, but the present invention shall not be

5 ~~restricted to the examples described below.~~

Examples 1 to 8 and Comparative Examples 1 to 7

The respective hair growth promotor compositions were prepared according to formulations shown in the
10 following Table 1 and Table 2. The blend unit is mass %, and the total amount is 100 mass % (the same shall apply to Example 9 and examples subsequent thereto).

The hair growth promotor compositions (samples to be tested falling in the scope of the present invention
15 (examples) and samples to be tested falling outside the scope of the present invention (comparative examples)) thus obtained were evaluated for a hair growth effect, use feeling and stability at a low temperature by the respective methods described below. The results thereof
20 are shown in Table 1 and Table 2.

Evaluation method of hair growth effect:

The volunteers of 64 adult men were divided at random into total 8 groups as a subject group and a
25 control group, in which one group consists of 8 men.

The test was carried out by applying adequate amounts of the compositions onto a frontal region through a parietal region twice a day, in the morning and at night. The administration period was 4 months.

- 5 ~~An improvement degree of the hairs at the time of~~ finishing the test (4 months) as compared with those before starting the test was judged according to the following five stage evaluation (notable improvement, middle improvement, light improvement, unchanged and
- 10 degraded) referring to observation of the photos.

Judgment criteria:

notable improvement: vellus hairs were scarcely observed and normalized

- middle improvement: vellus hairs were turned to terminal
- 15 hairs to a considerable extent

light improvement: vellus hairs were slightly turned to terminal hairs

unchanged: change was not observed at all on the quality of the hairs

- 20 degraded: the hairs were turned to vellus hairs.

The evaluation results of the hair growth effect shown in Table 1 and Table 2 were shown by a proportion (% , improvement rate) of the number of the persons having the effects of notable improvement ~ light improvement

25 after used respectively by 8 persons.

Evaluation method of use feeling:

The use feeling (average) in evaluating the hair growth-accelerating effect described above, that is, in
 5 applying the adequate amount onto a frontal region through a parietal region twice a day, in the morning and at night, was evaluated according to the following evaluation criteria.

Evaluation criteria:

10 ○: hairs are not dried up, and good use feeling is obtained

△: hairs are dried up a little, but use feeling is not dissatisfactory

×: hairs are dried up, and use feeling is
 15 dissatisfactory

Evaluation method of stability at low temperature:

A transparent glass bottle having a cap was filled with about 50 mL of the respective samples obtained and
 20 then stored at -10°C to evaluate deposition of crystals and the presence of precipitates with eyes after 4 weeks according to the following evaluation criteria.

Evaluation criteria:

○: crystals and precipitates are not observed

25 △: trace amounts of crystals and precipitates are

observed

X: crystals or precipitates are observed

Table 1

		Blend amount (%)							
		Example							
		1	2	3	4	5	6	7	8
Component (A)	Glycerin tridecanoate	2.0	2.0	2.0				2.0	
	Glycerin pentadecanoate				2.0	2.0	2.0		2.0
Component (B)	6-Benzylamino-Purine	0.5	0.5		0.5				0.5
	6-(4-Methylbenzyl-amino)purine			0.5		0.5	0.5	0.5	
Component (C)	Pentaglycerin monomyristate	0.5			1.0			0.05	
	Decaglycerin monomyristate		0.5			2.0			0.05
	Pentaglycerin monooleate			1.0			2.0		
Component (D)	Sorbitan monolaurate	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
99 % ethanol		Balance	Balance	Balance	Balance	Balance	Balance	Balance	Balance
Evaluation of hair growth effect		75	75	63	75	75	75	63	63
Evaluation of use feeling		○	○	○	○	○	○	△	△
Evaluation of stability at low temperature		○	○	○	○	○	○	○	○

Table 2

		Blend amount (%)						
		Comparative Example						
		1	2	3	4	5	6	7
Component (A)	Glycerin tridecanoate	2.0	2.0	2.0				
	Glycerin pentadecanoate				2.0	2.0	2.0	2.0
Component (B)	6-Benzylamino-purine	0.5	0.5					0.5
	6-(4-Methylbenzyl-amino)purine			0.5	0.5	0.5	0.5	
Component (C)	Pentaglycerin monomyristate		2.0					
	Decaglycerin monomyristate				2.0			
	Pentaglycerin monooleate							2.0
	Sorbitan monolaurate			3.0			1.0	
99 % ethanol		Balance	Balance	Balance	Balance	Balance	Balance	Balance
Evaluation of hair growth effect		75	75	63	75	75	75	63
Evaluation of use feeling		X	Δ	○	Δ	X	○	Δ
Evaluation of stability at low temperature		X	X	X	X	X	X	X

As apparent from the results shown in Table 1 and Table 2, it has become clear that in all of Examples 1 to 8 containing the prescribed amounts of the active ingredients (the component (A) to the component (D))

- 5 falling in the scope of the present invention, excellent characteristics are shown in the hair growth effect, the use feeling and the stability at low temperature and that in all of Comparative Examples 1 to 7 falling outside the scope of the present invention, excellent characteristics
10 are not shown in the use feeling and the stability at low temperature.

To specifically observe the comparative examples, Comparative Examples 1 and 5 are cases in which the component (A) and the component (B) are contained and
15 both of the component (C) and the component (D) are not contained, and Comparative Examples 2 to 4, 6 and 7 are cases in which the component (A) and the component (B) are contained and either one of the component (C) and the component (D) is contained. It can be found that the
20 effects of the present invention can not be exhibited in the above cases.

Examples 9 to 20

Next, examples in which the hair growth promotor
25 compositions of the present invention have been applied

to commercial products shall be shown. The hair growth promotor compositions in the respective examples described below were prepared according to the respective formulations by ordinary methods for preparing the

5 respective preparations.

The hair growth promotor compositions of various preparations (a hair growth promotor, a hair growth spray, a hair growth tonic and a hair growth lotion) prepared in Examples 9 to 20 described below were evaluated for a
 10 hair growth effect, use feeling and stability at low temperature in the same manners as in Examples 1 to 8 to find that the same excellent effects as in Example 1 were shown in all examples.

15 Example 9 (hair growth promotor)

Formulation

	Glycerin monopentadecanoate	3.0
	6-Benzylaminopurine	0.5
	DL- α -tocopheryl acetate	0.1
20	Coleus forskohlii root extract	0.5
	Glycerin	0.5
	Sorbitan coconut oil fatty acid ester	1.0
	Sucrose myristic acid ester	0.5
	Decaglycerin monomyristate	1.0
25	Lauryldimethylamino acetic acid betaine	0.3

	Amphoteric polymer*	0.2
	Ethyl oleate	2.0
	Succinic acid	0.1
	Fragrance#	0.5
5	Refined water	0.3
	99.5 % ethanol	balance

*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine-alkyl methacrylate copolymer

#: an A composition in Table 2 described in Japanese

10 Patent Application Laid-Open No. 113019/2003

Example 10 (hair growth promotor)

Formulation

	Glycerin monopentadecanoate	2.0
15	6-Benzylaminopurine	0.1
	β -glycyrrhetic acid	0.1
	Amphoteric polymer*	0.2
	Ethyl oleate	1.0
	Succinic acid	0.3
20	Sucrose lauric acid ester	0.5
	Sorbitan monolaurate	0.5
	Glycerin	0.8
	Decaglycerin monomyristate	0.5
	L-menthol	0.1
25	DL- α -tocopheryl acetate	0.1

	Refined water	0.3
	Fragrance#	0.5
	99.5 % ethanol	balance
5	Prepared fluid	
	Formulated concentrate described above	80 %
	LPG	20 %
	*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine-alkyl methacrylate copolymer	
10	#: a B composition in Table 2 described in Japanese Patent Application Laid-Open No. 113019/2003	
	Example 11 (hair growth tonic)	
	Formulation	
15	Glycerin monopentadecanoate	2.0
	6-Benzylaminopurine	0.05
	POE (8 mole) oleyl ether	1.5
	Sorbitan monomyristate	3.0
	Pentaglycerin monolaurate	0.3
20	L-menthol	0.1
	Hinokitiol	0.3
	Methylparaben	0.1
	Fragrance#	0.3
	Refined water	0.3
25	99.5 % ethanol	balance

#: a C composition in Table 2 described in Japanese
Patent Application Laid-Open No. 113019/2003

Example 12 (hair growth lotion)

5	Formulation	
	Glycerin monopentadecanoate	1.0
	6-(4-Methylbenzylamino)purine	0.5
	Natural vitamin E	0.5
	Sucrose myristic acid ester	0.5
10	POE (40) hardened castor oil	0.5
	Pentaglycerin monooleate	0.5
	Sorbitan monomyristate	0.8
	Citric acid	0.1
	L-menthol	0.1
15	Fragrance#	0.5
	Refined water	0.3
	99.5 % ethanol	balance

#: a D composition in Table 2 described in Japanese
Patent Application Laid-Open No. 113019/2003

20

Example 13 (hair growth promotor)

Formulation

	Glycerin monopentadecanoate	3.0
	6-Benzylaminopurine	0.4
25	D-pantothenyl alcohol	0.4

	β -glycyrrhetic acid	0.1
	Pyrocton olamin	0.1
	Gallic acid 3,5-diglucoside	0.1
	Pentaglycerin monooleate	0.8
5	Coleus extract	1.2
	Stearyltrimethylammonium chloride	0.8
	Sorbitan monolaurate	0.5
	Lauryldimethylamino acetic acid betaine	0.3
	Amphoteric polymer*	0.2
10	Ethyl oleate	2.0
	Succinic acid	0.1
	Fragrance#	0.5
	Refined water	0.3
	99.5 % ethanol	balance
15	*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine-alkyl methacrylate copolymer	
	#: an A composition in Table 2 described in Japanese Patent Application Laid-Open No. 113019/2003	
20	Example 14 (hair growth promotor)	
	Formulation	
	Glycerin monopentadecanoate	3.0
	6-Benzylaminopurine	0.4
	Nicotinamide	0.4
25	β -glycyrrhetic acid	0.1

	Pyrocton olamin	0.1
	Gallic acid-3,5-diglucoside	0.1
	Methyl salicylate	0.5
	Sorbitan monooleate	3.0
5	Decaglycerin monomyristate	0.5
	Stearyltrimethylammonium chloride	0.4
	Polyoxyethylene oleyl ether	0.5
	Lauryldimethylamino acetic acid betaine	0.3
	Amphoteric polymer*	0.2
10	Polyethylene glycol	2.0
	L-menthol	0.1
	Fragrance#	0.2
	Succinic acid	0.1
	Ethanol	balance
15	*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine-alkyl methacrylate copolymer	
	#: a C composition in Table 2 described in Japanese Patent Application Laid-Open No. 113019/2003	
20	Example 15 (hair growth promotor)	
	Formulation	
	Glycerin monopentadecanoate	3.0
	6-Benzylaminopurine	0.4
	Nicotinamide	0.4
25	Tocopheryl acetate	0.1

	Pyrocton olamin	0.1
	Gallic acid-3,5-diglucoside	0.05
	Propylene glycol	6.0
	Stearyltrimethylammonium chloride	0.6
5	Decaglycerin monomyristate	0.5
	Sorbitan monolaurate	5.0
	Polyoxyethylene oleyl ether	0.5
	Polyethylene glycol	2.0
	L-menthol	0.2
10	Nylon powder*1	1.5
	Fragrance#	0.2
	Ethanol	balance

*1: nylon 12, average particle diameter: about 10 μm

#: a B composition in Table 2 described in Japanese

15 Patent Application Laid-Open No. 113019/2003

Example 16 (hair growth promotor)

Formulation

	6-Benzylaminopurine	0.5
20	Glycerin monopentadecanoate	3.0
	Gallic acid-3,5-diglucoside	0.3
	Tocopheryl acetate	0.1
	Sorbitan palmitate	2.0
	β -glycyrrhetic acid	0.5
25	Sorbitan coconut oil fatty acid ester	0.5

	Stearyltrimethylammonium chloride	0.3
	Sucrose fatty acid ester	0.3
	Decaglycerin monomyristate	0.6
	Amphoteric polymer*	0.02
5	Ethyl oleate	1.5
	Pantothenyl ethyl ether	1.0
	D-pantothenyl alcohol	1.0
	Coleus forskohlii root extract	2.0
	Conc. glycerin	1.0
10	Isodonis Japonicus extract	1.0
	Succinic acid	0.1
	Benzalkonium chloride	0.02
	L-menthol	0.1
	Ethanol	balance
15	*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine-alkyl methacrylate copolymer	

Example 17 (hair growth promotor)

Formulation

20	6-Benzylaminopurine	0.5
	Glycerin monopentadecanoate	3.0
	Tocopheryl acetate	0.1
	Sorbitan coconut oil fatty acid ester	2.0
	Sucrose fatty acid ester	0.3
25	Decaglycerin monomyristate	0.6

	Amphoteric polymer*	0.5
	Ethyl oleate	1.5
	Coleus·forskohlii root extract	1.5
	Glycerin	1.0
5	-Glycerin monomyristate	0.05
	Pentadecanoic acid	0.01
	Myristic acid	0.01
	Glycerin dipentadecanoate	0.05
	Succinic acid	0.1
10	L-menthol	0.1
	Fragrance#	0.2
	Ethanol	balance

*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine·alkyl methacrylate copolymer

15 #: a C composition in Table 2 described in Japanese Patent Application Laid-Open No. 113019/2003

Example 18 (hair growth promotor)

Formulation

20	6-Benzylaminopurine	0.5
	Glycerin monopentadecanoate	2.0
	Tocopheryl acetate	0.1
	β -glycyrrhetic acid	0.1
	Sorbitan palmitate	2.0
25	Sorbitan coconut oil fatty acid ester	0.5

	Sucrose fatty acid ester	0.3
	Decaglycerin monomyristate	0.6
	Amphoteric polymer*	0.02
	Ethyl oleate	1.5
5	-Pantothenyl ethyl ether	-1.0
	Coleus forskohlii root extract	2.0
	Pentadecanoic acid	0.05
	Palmitic acid	0.01
	Glycerin monopalmitate	0.02
10	Glycerin dipentadecanoate	0.05
	Glycerin	0.03
	Isodonis Japonicus extract	1.0
	Succinic acid	0.1
	L-menthol	0.2
15	Fragrance#	0.2
	Ethanol	balance

*: N-methacryloylethyl-N,N-dimethylammonium- α -N-methylcarboxybetaine-alkyl methacrylate copolymer

#: a D composition in Table 2 described in Japanese

20 Patent Application Laid-Open No. 113019/2003

Example 19 (hair growth promotor)

Formulation

	6-Benzylaminopurine	0.5
25	Glycerin monopentadecanoate	3.0

	Tocopheryl acetate	0.1
	Gallic acid-3,5-diglucoside	0.3
	Sorbitan coconut oil fatty acid ester	2.0
	Sucrose fatty acid ester	0.3
5	Ethyl oleate	1.5
	Decaglycerin monomyristate	0.6
	Amphoteric polymer*	0.5
	Stearyltrimethylammonium chloride	0.3
	Coleus forskohlii root extract	1.5
10	Glycerin	1.0
	Pentadecanoic acid	0.01
	Myristic acid	0.01
	Glycerin monomyristate	0.05
	Glycerin dipentadecanoate	0.05
15	Succinic acid	0.1
	L-menthol	0.1
	Fragrance#	0.2
	Ethanol	balance

*: N-methacryloylethyl-N,N-dimethylammonium- α -N-

20 methylcarboxybetaine-alkyl methacrylate copolymer

#: an E composition in Table 2 described in Japanese

Patent Application Laid-Open No. 113019/2003

Example 20 (hair growth promotor spray)

25 Formulation

	6-Benzylaminopurine	0.5
	Glycerin monopentadecanoate	3.0
	Tocopheryl acetate	0.1
	Gallic acid-3,5-diglucoside	0.3
5	—Sorbitan coconut oil fatty acid ester	2.0
	Sucrose fatty acid ester	0.3
	Ethyl oleate	1.5
	Decaglycerin monomyristate	0.6
	Amphoteric polymer*	0.5
10	Stearyltrimethylammonium chloride	0.3
	Coleus forskohlii root extract	1.5
	Glycerin	1.0
	Pentadecanoic acid	0.01
	Myristic acid	0.01
15	Glycerin monomyristate	0.05
	Glycerin dipentadecanoate	0.05
	Succinic acid	0.1
	L-menthol	0.1
	Fragrance#	0.2
20	Ethanol	balance
	Prepared fluid	
	Formulated concentrate described above	80 %
	LPG	20 %
25	*: N-methacryloylethyl-N,N-dimethylammonium- α -N-	

methylcarboxybetaine·alkyl methacrylate copolymer

#: an E composition in Table 2 described in Japanese
Patent Application Laid-Open No. 113019/2003

5 - Industrial Applicability

The hair growth promotor composition of the present invention thus constituted exhibits an excellent hair growth effect by applying externally onto a scalp and is excellent in stability at low temperature, and it provides good use feeling having no stickiness, so that the hair growth promotor composition having excellent effects which have not so far been shown is provided.